

UT Southwestern Department of Radiology

Protocol Name: Ankle/Foot CT With IV Contrast

Orderable Name: CT LOWER EXTREMITY LEFT W IV CONTRAST
CT LOWER EXTREMITY RIGHT W IV CONTRAST

Adult Only

Epic Button: Ankle/Foot + IV

CTDIvol < 50 mGy

Indications: Suspected infection, inflammation, tumor

Acquisitions: 1

Active Protocol

<p>Oral Contrast: None</p>	<p>IV Contrast: Link to Contrast Information</p> <p>Rate (mL/sec): 3</p> <p>Volume (mL): 75</p> <p>IV Access: Power injection: 20g or larger strongly preferred (if 22g use reduce rate to 2.5 mL/sec)</p> <p>Notes:</p>	<p>Other Contrast: None</p>	<p>Airway</p> <hr/> <p>Other Notes *Place a marker at the site of most concern. Dual energy scanner required at CUH & OPB. Use Right/Left orderable based on protocol or side indicated in reason for exam. Metal (FOV):140 Kvp,align beam along metal.</p>
-----------------------------------	---	------------------------------------	---

Last Change: 10/6/2022

Last Review: 1/24/2022

Links: [General Statements](#)

Special Instructions	Use 5mm cor/sag if large patient or metal in FOV.	Do not repeat CT scan, recon soft tissue from 1st acquisition, send soft tissue kernel volume to TeraRecon Use 5mm cor/sag if large patient or metal in FOV.
Acq # / Series Name	1 60 Sec Delayed	N/A 60 Sec Delayed
Phase Timing	60 seconds	N/A
Acquisition Protocol		Recon Only
Coverage	Distal tibia thru toes	Same
FOV	Whole ankle and foot	Same
Algorithm	Bone	Soft Tissue
Primary Axial Recon	3 mm	3 mm, volume
Other Recons	3 mm coronal and sagittal	4 mm coronal and sagittal
MIP Recons		
*Dual Energy Philips	VNC, Gout maps (cor/sag), BM edema, SBI, mono E 100, mono E 120	
*Dual Energy Siemens	VNC, Gout maps (cor/sag), BM edema, low/high kVp, mono E 100, mono E 120	



* When dual energy or spectral CT is used