UT Southwestern Department of Radiology

Protocol Name: CT Head Without and With Orderable Name: CT BRAIN W AND WO IV CONTRAST

Epic Button: Head W/O & W/ CTDIvol < 80 mGy

Indications: Infection, malignancy, AVM, aneurysm, stroke, seizures.

Acquisitions: 2

Active Protocol

Adult Only

Oral Contrast: None	IV Contrast: Link to Contrast Information	Other Contrast: None	Airway
	Rate (ml/sec): 3		
	Volume (ml): 70		
	IV Access:		Other Notes
	Power injection: 20g or larger strongly preferred (if 22g use		No habitus specific kVp adjustment.
	reduce rate to 2.5 mL/sec)		Utilize Dual Energy when available
	Notes:		
	Saline Test Injection: 10 sec @ injection rate		

Last Change: 1/24	/2024	Last Review: 1/8/2025	Links:				
Special Instructions			Do not repeat CT scan, recon soft tissue from 1st acquisition Send soft tissue volume to PACS. Send low/hi kVp to SyngoVia.				
Acq # / Series Name	1	Noncontrast		N/A	Noncontrast	2	Contrast
Phase Timing	Phase Timing		ı	N/A		60 seconds	
Acquisition Protocol	cquisition Protocol Neuro Head		F	Recon Only		Neuro Head	
Coverage Base of skull through vertex		Ş	Same				
FOV	FOV 200 mm		9	Same		200 mm	
Algorithm Bone		9	Soft Tissue		Soft Tissue		
Axial Recons 5 mm			5 mm, 1 mm (volume)		5 mm, 1 mm (volume)		
Other Planar Recons			3	3 mm coronal and sagittal		3 mm coronal and sagittal	
MIP Recons							
†DECT Philips							
†DECT Siemens			-	1mm SF	op.		
†PC-CT Siemens							

